

Literature:

Clinical studies listed below demonstrate usage of Microdialysis monitoring in the gastrointestinal tract following surgery.

Intraperitoneal microdialysis in the postoperative surveillance of infants undergoing surgery for congenital abdominal wall defect: A pilot study.

J Pediatr Surg. 2015 Mar 10.

Risby K, Ellebæk MB, Jakobsen MS, Husby S, Qvist N.

Mediastinal microdialysis in the diagnosis of early anastomotic leakage after resection for cancer of the esophagus and gastroesophageal junction.

Am J Surg. 2014 Jan 16.

Ellebæk M, Qvist N, Fristrup C, Mortensen MB.

Identification of anastomotic leakage after colorectal surgery using microdialysis of the peritoneal cavity.

Tech Coloproctol. 2013 Apr 30

Daams F, Wu Z, Cakir H, Karsten TM, Lange JF.

Intra-peritoneal Microdialysis and Intra-abdominal Pressure after Endovascular Repair of Ruptured Aortic Aneurysms.

Eur J Vasc Endovasc Surg. 2013 Mar 26

Hörner TM, Skoog P, Norgren L, Magnuson A, Berggren L, Jansson K, Larzon T.

Intraperitoneal glycerol levels and lactate/pyruvate ratio: early markers of postoperative complications.

Scand J Gastroenterol. 2011 Mar 28.

Hörner TM, Norgren L, Jansson K.

Intraperitoneal microdialysis in the postoperative surveillance after surgery for necrotizing enterocolitis: a preliminary report.

Pedersen ME, Dahl M, Qvist N. J Pediatr Surg. 2011 Feb

Peritoneal microdialysis. Early diagnosis of anastomotic leakage after low anterior resection for rectosigmoid cancer.

Scand J Surg. 2009

Ellebæk Pedersen M, Qvist N, Bisgaard C, Kelly U, Bernhard A, Møller Pedersen S.


Intraperitoneal microdialysis: Postoperative monitoring of splanchnic ischemia by measurements of the lactate pyruvate ratio.

Crit Care Med. 2006 Oct

Jansson M, Strand I, Jansson K.

Contact information

Principle of Microdialysis



M Dialysis AB

M Dialysis is the leading company devoted to the development, manufacturing and marketing of the Microdialysis technique.

The head office is located in Stockholm, Sweden, with a subsidiary in Boston MA, USA. M Dialysis has distributors across the globe, responsible for local sales, service and support.



M Dialysis AB, Box 5049, SE-121 05 Stockholm, Sweden,
Tel: +46 8 470 10 20, Fax: +46-8-470 10 55
E-mail: info@mdialysis.se, www.mdialysis.com

Microdialysis

Gastrointestinal Surgery

Ref: No. 8050156A 2017-08-21

Distributor

 dialysis

Microdialysis in Gastrointestinal Surgery

Gastrointestinal surgery is associated with a high risk of post-operative complications such as anastomosis leakages and sepsis. Early diagnosis of complications and intervention is critical to improve outcome.

Intraperitoneal Microdialysis is a unique technique for early detection of complications following surgery. The technology is minimally-invasive, easy to handle, and can be used for several days. The method is performed by inserting a Microdialysis catheter into the intraperitoneal space during open surgery.

The catheter is perfused with a sterile isotonic solution via a small pump attached to its inlet lumen. When inserted in the intraperitoneal space small substances diffuse through the semi-permeable Microdialysis membrane into the perfusion fluid. This fluid, now known as dialysate, moves through the outlet lumen and into a collection microvial. The Microvials are exchanged at regular intervals and analyzed immediately using the ISCUS^{flex} Microdialysis Analyzer.

The metabolite values in the collected sample provide a picture of the local metabolism and Microdialysis is a safe and reliable monitoring for surveillance of for example an anastomosis after colorectal surgery.

Early detection of metabolic changes following Gastro Intestinal Surgery




The 62 Gastro Intestinal Microdialysis Catheter is placed free floating in the intraperitoneal space close to the anastomosis during open surgery.

Microdialysis sampling

Microdialysis sampling is carried out by placing the sterile CE-certified 62 Gastrointestinal Microdialysis catheter in the Intraperitoneal space close to the anastomosis after colorectal surgery.

62 Gastrointestinal Microdialysis Catheter



- For Implantation in the intraperitoneal space during open surgery
- Sterile, single use

106 Microdialysis Pump



The 106 Microdialysis Pump is dedicated for the perfusion of Microdialysis catheters with sterile isotonic perfusion fluid. It is handy and battery driven. The operating status is indicated by LED's and it operates at a fixed flow rate of 0.3 µl/min.

The Microdialysis monitoring system is light and portable and is not disturbing nursing actions or movement of the patient.

ISCUS^{flex} Microdialysis Analyzer

The ISCUS^{flex} Microdialysis Analyzer is specially designed for the handling of small Microdialysis sample volumes.

The system is unique for monitoring metabolic changes in tissues, intraperitoneal fluid and organs during surgery, in intensive care and normal ward.

Biochemical markers:

- Glucose
- Glutamate
- Lactate
- Glycerol
- Pyruvate
- Urea
- LP-ratio



The ISCUS^{flex} Microdialysis Analyzer is easily operated by medical professionals. Data is displayed as trend curves for easy and fast interpretation.